

CORRECTION

Prevalence of Obesity and its Association with Blood Pressure, Serum Lipids and Selected Lifestyles in a Puerto Rican Population of Adolescents. Venegas HL, Pérez CM, Suarez EL, Guzmán M. (PR Health Sci J, June 2003; 22:137-143). In Table 7, p. 141, the names of the specific serum lipids were left out. Please see below the corrected Table 7. We regret the error.

Table 7. Prevalence Odds Ratio (POR) between Obesity* and Serum Lipids in a Sample of Volunteers in a Junior-High School in San Juan, Puerto Rico, 1999-2000 (n=26).

| Serum lipid | Levels Above Median | Levels Below Median | POR (95% CI) | p-value† |
|----------------------------------|---------------------|---------------------|--------------|----------|
| Total cholesterol (mg/dl) | | | | |
| Obese | 8 | 3 | 5.3 | 0.55 |
| Non-obese | 5 | 10 | (0.76, 43.0) | |
| HDL-C (mg/dl) | | | | |
| Obese | 7 | 4 | 2.0 | 0.32 |
| Non-obese | 7 | 8 | (0.31, 13.4) | |
| LDL-C (mg/dl) | | | | |
| Obese | 7 | 4 | 2.6 | 0.21 |
| Non-obese | 6 | 9 | (0.41, 17.8) | |
| Triglycerides (mg/dl) | | | | |
| Obese | 6 | 5 | 1.37 | 0.50 |
| Non-obese | 7 | 8 | (0.22, 8.6) | |

* Obese group included level I and level II obesity

† Fisher's exact test